

NADA Number: 033-157	
Trade Name	Spectam® Scour Halt Spectam® Scour Halt®
Sponsor	Teva Animal Health, Inc.
Ingredients	Spectinomycin Hydrochloride Pentahydrate
Species	Swine, neonate or suckling
Routes of Administration	Per Os
Dose Form	Liquid (solution)
Drug Form	Liquid (solution)
Dispensing Status	OTC
Dosage Amount, Indications & Limitations	<p>520.2123c Spectinomycin dihydrochloride pentahydrate solution.</p> <p>Specifications: Each milliliter of solution contains 50 milligrams spectinomycin activity.</p> <p>Conditions of use:</p> <p>Swine (under 4 weeks and under 15 pounds).</p> <p>Amount: Administer 5 milligrams per pound of body weight orally twice daily for 3 to 5 days.</p> <p>Indications: It is used for the treatment and control of infectious bacterial enteritis (white scours) associated with E. coli in pigs under 4 weeks of age.</p> <p>Limitations: Do not administer to pigs over 15 pounds body weight or over 4 weeks of age. Do not administer within 21 days of slaughter.</p>
Tolerances	<p>A tolerance of 0.1 part per million (ppm) for negligible residues of spectinomycin in uncooked edible tissues of chickens and turkeys is established.</p> <p>A tolerance of 4 ppm for parent spectinomycin (marker residue) in kidney (target tissue) and 0.4 ppm for parent spectinomycin in cattle muscle in cattle is established.</p>